arXiv:1207.1483v1 [cond-mat.quant-gas] 5 Jul 2012

Fast generation of spin-squeezed states in bosonic Josephson junctions

B. Juliá-Díaz[1], E. Torrontegui[2], J. Martorell[3], J. G. Muga[2,4], and A. Polls[3]
[1] *ICFO-Institut de Ciències Fotòniques, Parc Mediterrani de la Tecnologia, 08860 Barcelona, Spain*
[2] *Departamento de Química-Física, UPV-EHU, Apartado 644, 48080 Bilbao, Spain*
[3] *Deptartament d'Estructura i Constituents de la Matèria,*
Facultat de Física, U. Barcelona, 08028 Barcelona, Spain and
[4] *Department of Physics, Shanghai University, 200444 Shanghai, Peoples Republic of China*

We describe methods for fast production of highly coherent-spin-squeezed many-body states in bosonic Josephson junctions (BJJs). We start from the known mapping of the two-site Bose-Hubbard (BH) Hamiltonian to that of a single effective particle evolving according to a Schrödinger-like equation in Fock space. Since, for repulsive interactions, the effective potential in Fock space is nearly parabolic, we extend recently derived protocols for shortcuts to adiabatic evolution in harmonic potentials to the many-body BH Hamiltonian. The best scaling of the squeezing parameter for large number of atoms N is $\xi_S^2 \sim 1/N$.

Recent experimental results from the group of M. Oberthaler have provided evidence of the generation of spin-squeezed states in bosonic Josephson junctions [1–4]. Two different setups with ultracold ^{87}Rb bosons have been explored: a) a cloud trapped in an external double-well potential [1, 5] (external BJJ) and b) a cloud of atoms in two different hyperfine states trapped in a single harmonic potential with a linear coupling between the two internal states [4] (internal BJJ). In both cases, the atomic-clouds are well described by the two-site Bose-Hubbard model [5–7], a Lipkin-Meshkov-Glick-like (LMG) Hamiltonian [8]. As previously pointed out [9], this Hamiltonian can produce largely squeezed spin states and, correspondingly, large sets of entangled pseudo-spins [3, 10, 11].

Simultaneously, and due to the need to control and implement quantum resources, there has been an increasing interest in developing fast protocols to shortcut adiabatic following [12, 13]. The key purpose is to engineer procedures to drive, in a finite time, a system from some initial state to a final state that could be reached with an adiabatic, slow process. The different protocols may in addition aim at a) minimizing the transient energy excitations, b) reducing the sensitivity to noise, or c) minimizing other variables of interest [14]. It should be noted that, in general, the initial and final states are not necessarily required to be the ground states (GSs) of the system. In this work, however, we concentrate in transitions between ground states corresponding to different values of the model parameters which can be controlled externally. The proposed method is designed so that the desired final state is produced as a stationary eigenstate of the Hamiltonian, with no need to stop or freeze the dynamics. Analytical formulas to perform this type of processes exist for the harmonic oscillator [12]. We show that they can be extended to many-body states described by the BH Hamiltonian. That allows us to compute the dependence of the attainable coherent spin squeezing in the system[1] on the number of atoms, N, the atom-atom interactions, U, and the linear coupling J. To do so, we benefit from the explicit mapping, in the large N limit, between the two-site Bose-Hubbard model and an effective single particle system described by a Schrödinger-like equation [15–17] (in the more general context of the LMG model, this connection was established earlier [18, 19], albeit with no known relation at the time with ultracold atom physics).

The dynamics of a bosonic Josephson junction can be well described by a quantized two-mode model [5–7]. For the external BJJ case [1, 5], or by construction in the internal BJJ case, the system of ultra-cold bosons is well modeled by the Bose-Hubbard Hamiltonian $\hbar\mathcal{H}_{\rm BH}$

$$\mathcal{H}_{\rm BH} = -2J\hat{J}_x + U\hat{J}_z^2\,, \tag{1}$$

where the pseudo-angular momentum operator $\hat{\mathbf{J}} \equiv \{\hat{J}_x, \hat{J}_y, \hat{J}_z\}$ is defined as, $\hat{J}_x = (1/2)(\hat{a}_1^\dagger\hat{a}_2 + \hat{a}_2^\dagger\hat{a}_1)$, $\hat{J}_y = (1/2i)(\hat{a}_1^\dagger\hat{a}_2 - \hat{a}_2^\dagger\hat{a}_1)$, and $\hat{J}_z = (1/2)(\hat{a}_1^\dagger\hat{a}_1 - \hat{a}_2^\dagger\hat{a}_2)$. $\hat{a}_j^\dagger$ creates a boson in site j, $[\hat{a}_i, \hat{a}_j^\dagger] = \delta_{i,j}$. J is the hopping strength, taken positive, and U is the non-linear coupling strength proportional to the atom-atom s-wave scattering length[2]. In this work we consider repulsive interactions, $U > 0$. For internal BJJ, the inter-species s-wave scattering length in ^{87}Rb atoms can be varied by applying an external magnetic field thanks to a well characterized Feshbach resonance at $B = 9.1$ G, as done in Ref. [4] for the setup that we are considering, thus permitting precise control over the U. In this work, we use a time dependent $U(t)$, keeping J and N fixed during the time evolution. To do so one needs to track the resonance of the linear coupling (responsible for the hopping term in the Hamiltonian) which is altered due to the second order Zeeman shift when B is varied. Albeit technically challenging, this is within reach of the current experimental setups [20].

---

[1] The use of the spin squeezing as a witness of $N-$particle entanglement is established in Ref. [11]. In this letter we focus only in the characterization of the squeezing.

[2] In the internal BJJ, U is proportional to $a_{1,1} + a_{2,2} - 2a_{1,2}$, with $a_{1,1}$ and $a_{2,2}$ the intra-species scattering lengths and $a_{1,2}$ the inter-species one [4].

The time dependent Schrödinger equation (TDSE) is written as, $\imath\partial_t|\Psi\rangle = \mathcal{H}_{\rm BH}|\Psi\rangle$. For a given N, an appropriate many-body basis for this bosonic system is the Fock basis, $\{|m_z = (N_1 - N_2)/2\rangle\}$, with $m_z = -N/2, \ldots, N/2$. A general many-body state, $|\Psi\rangle$, can then be written as $|\Psi\rangle = \sum_{m_z=-N/2}^{N/2} c_{m_z}|m_z\rangle$.

As customary, the number squeezing parameter is defined as [2, 21, 22], $\xi_N^2(t) = \Delta\hat{J}_z^2/(\Delta\hat{J}_z^2)_{\rm bin}$ where $\Delta\hat{J}_z^2 \equiv \langle\hat{J}_z^2\rangle - \langle\hat{J}_z\rangle^2$ and $(\Delta\hat{J}_z^2)_{\rm bin} = N/4$ in the binomial case. The many-body state is said to be number-squeezed if $\xi_N < 1$ [9]. The coherent spin-squeezing parameter is defined as [9, 10, 21] $\xi_S^2 = N(\Delta\hat{J}_z^2)/\langle\hat{J}_x\rangle^2 = \xi_N^2/\alpha^2$, where the phase coherence of the many-body state is $\alpha(t) = \langle\Psi(t)|2\hat{J}_x/N|\Psi(t)\rangle$. ξ_S takes into account the delicate compromise between improvements in number-squeezing and loss of coherence. States with $\xi_S < 1$ have been proposed to be used in a new Ramsey type atom interferometer with an increased phase precision compared to the coherent spin state. This gain in precision can be directly related to entanglement in the system [10].

As detailed in Refs. [15–17] a systematic expansion of $\mathcal{H}_N \equiv \mathcal{H}_{\rm BH}/(NJ)$ in powers of $h \equiv 1/N$ gives in the semiclassical limit, $N >> 1$, the pseudo-Schrödinger-like equation,

$$\mathcal{H}_N(z)\psi(z) \equiv -2h^2\partial_z\sqrt{1-z^2}\,\partial_z\psi(z) + \mathcal{V}(z)\psi(z) \quad (2)$$

where $\mathcal{V}(z) = -\sqrt{1-z^2} + (1/2)\Lambda z^2$, $z = m_z/(N/2)$, $\Lambda = NU/(2J)$ and $\psi(z) = \sqrt{N/2}\,c_{m_z}$, normalized as $\int_{-1}^{1} dz|\psi(z)|^2 = 1$. The corresponding TDSE writes as $ih\partial_t\psi(z,t) = \mathcal{H}_N\psi(z,t)$ with time measured in units of $1/J$. In these units the Rabi time is defined as $t_{\rm Rabi} = \pi/J$.

As explained in [17], and previously noted by other authors [15, 23], for repulsive interactions the potential in Fock space, $\mathcal{V}(z)$, is to a very good approximation a harmonic oscillator: Neglecting the z dependence of the effective mass term and expanding $\sqrt{1-z^2} \simeq 1 - z^2/2$ in $\mathcal{V}(z)$: Eq. (2) reduces to

$$\mathcal{H}_N \simeq -2h^2\partial_z^2 + \frac{1}{8}\omega^2 z^2\,, \quad (3)$$

with $\omega = 2\sqrt{1+\Lambda}$.

We have checked that all the expectation values of $\hat{J}_i$ and $\hat{J}_i^2$, $i = x, y, z$ computed by directly solving the TDSE for the BH Hamiltonian, and the corresponding ones using the solution Eq. (2) and the explicit expressions given in Ref. [17] agree almost perfectly in all calculations reported in this letter. The only minor discrepancy is explicitly shown in Fig. 3.



FIG. 1. (color online) Time evolution of the coherence $\alpha(t)$ and the number squeezing ξ_N^2 of the system computed using the fast adiabatic evolution of Eq. (4) (solid) compared to the case of linear ramping of $\Lambda(t)$ (dashed). The corresponding instantaneous adiabatic results for both the fast adiabatic case and a linear ramping are given in dotted and dot-dashed, respectively. $\Lambda(0) = 0$, $\Lambda(t_{\rm Rabi}) = 20$, and $N = 100$.

Fast shortcut to adiabatic evolution

Our method for fast adiabatic-like preparation of a given ground state requires control of the atom-atom interaction at time scales of fractions of the Rabi time. In the parabolic approximation discussed above the protocol developed in Ref. [12] has a direct translation into our variables. We need to impose the following time variation of Λ [3], $\Lambda(t) = \omega^2(t)/4 - 1$, where $\omega^2(t)$ obeys the Ermakov equation,

$$\ddot{b}(t)/J^2 + \omega^2(t)b(t) = \omega_0^2/b^3(t)\,, \quad (4)$$

with time in units of $1/J$ as explained above. The polynomial ansatz for $b(t)$ from Ref. [12] takes the explicit analytic form $b(t) = 6(r-1)s^5 - 15(r-1)s^4 + 10(r-1)s^3 + 1$, $s = t/t_f$, and $r = \sqrt{\omega_0/\omega_f}$ where $\omega_0 = 2\sqrt{1+\Lambda_0}$ and $\omega_f = 2\sqrt{1+\Lambda_f}$.

Our calculations confirm that this procedure, although not exact, is extremely accurate in the many-body simulations despite the fact that it was derived for the single particle problem in a harmonic oscillator Hamiltonian. This is due to the validity of the explained mapping of

---

[3] Note that the derivations in the previous section imply a fixed J and N, thus the variation in Λ has to be due to a variation in U.



FIG. 2. (color online) Considered $\Lambda(t)$ (upper panel): three final times are considered, $t_F = 0.2, 1$ and $5\,t_{\rm Rabi}$. Fidelity (middle panel) when $t_F = 0.2\,t_{\rm Rabi}$, and coherent spin squeezing ξ_S^2 (lower panel) as a function of t using the fast-adiabatic process described in the text. The adiabatic ξ_S^2, corresponding to the instantaneous ground state for each Λ, is shown in dashed lines in the lower panel. The initial and final Λ are 0 and 200. $N = 100$

the BH to the single particle pseudo-Schrödinger equation, which turns out to be well approximated by a harmonic oscillator potential. This harmonic approximation improves as N is increased because the spread of the GS wave function $\sim 1/\sqrt{N}$, thus exploring only the very central part of $\mathcal{V}(z)$ in $z \in [-1, 1]$. Exceptions appear in the ultra-fast preparation as discussed later.

Let's start by comparing the fast protocol to shortcut (SC) the adiabatic following presented above to the case of a linear ramping $\Lambda_l(t) = \Lambda(0) + (t/t_F)(\Lambda(t_F) - \Lambda(0))$. In this case we consider a relatively long final time of one full Rabi period $t_{\rm Rabi} = \pi/J$, see Fig. 1. For both $\Lambda(t)$ we present the results of the corresponding instantaneous ground state of the many-body system for comparison. First, we note that the SC does produce a final result which has similar coherence α and coherent number squeezing ξ_N^2 as the corresponding adiabatically evolved case. In contrast, the linearly ramped system deviates notably from the adiabatically evolved state in the final time. As we are considering fairly long times, the SC is not found to deviate appreciably from the instantaneous adiabatic following of the state, except for a short period at early times. The transient non-adiabaticity will be shown to increase as we require shorter t_F's.

The fidelity between the evolved many-body state $|\Psi(t)\rangle$ and the corresponding instantaneous ground-state, $|\langle\Psi(t)|\Psi_{\rm GS}\rangle|$, is plotted in Fig 2 and is seen to be extremely close to 1 at the final time for all considered t_F =0.2, 1 and 5 $t_{\rm Rabi}$. In the figure we compare the full many-body evolution to the corresponding instantaneous ground state for different t_F. As mentioned above, for short times, the fast-adiabatic passage produces intermediate many-body states which depart from the instantaneous ones, as seen clearly in the drop of the fidelity for very short times in the $t_F = 0.2\,t_{\rm Rabi}$ case.



FIG. 3. (color online) Considered $\Lambda(t)$ (upper panel), fidelity (middle panel) and coherent spin squeezing ξ_S^2 (lower panel) as a function of t for the exact Bose-Hubbard calculation (solid) and for the Schrödinger-like equation (2). The final time t_F is 0.05 $t_{\rm Rabi}$. The initial and final Λ are 0 and 500. $N = 100$. The inset zooms on the times close to t_F.

Ultra-short adiabatic production of squeezed states

As described in Ref. [12] for a particle in a harmonic oscillator and ultrafast adiabatic evolution, the method presented requires a certain period of the evolution where the trapping potential is actually reversed into an expelling potential (an inverted parabola). Through the previously explained mapping, this corresponds in our many-body BH problem to a period of time where the atom-atom interaction is switched from repulsive to attractive, $\Lambda < 0$. For attractive interactions the potential $\mathcal{V}(z)$ becomes a double-well potential in Fock space and the parabolic approximation of Eq. (3) does only hold for central regions in z [17]. Thus, one can expect that the present method should still work only when the wavefunction $\psi(z, t)$ does not spread in z significantly during the time where the atom-atom interactions are attractive. As seen in Fig. 3 this is indeed the case. We consider $t_F = 0.05\,t_{\rm Rabi}$, $\Lambda(0) = 0$ and $\Lambda(t_F) = 500$. Then the required $\Lambda(t)$ (upper panel) becomes negative for $0.025 \lesssim t/t_{\rm Rabi} \lesssim 0.04$. The method is seen to work accurately, producing a fidelity ~ 1 at the final time. The intermediate evolution is however highly diabatic,



FIG. 4. (color online) Coherent spin squeezing attainable using a simple "Bang-bang" method as described in the text (lower panel). The fidelity between the evolved state and its corresponding instantaneous evolution is given in the upper panel. $\Lambda(0) = 0$ in all cases, while $\Lambda(t_{\text{bang}}) = 5$ (dashed), 20 (dotted), and 200 (solid). $N = 100$.

with close to zero fidelity between the evolved state and its corresponding adiabatic one, during the $\Lambda(t) < 0$ evolution. The correspondence between the many-body BH and the single particle problem is made through the model of Eq. (2), which, as in all the other cases presented agrees well with the BH results.

Another set of protocols known to produce fast-adiabatic evolution in harmonic oscillator Hamiltonians are the so-called "bang-bang" methods. The translation of one of the simplest versions of these methods to the present problem requires the following steps assuming that we start as before from an initial ground state corresponding to $\Lambda(0) = \Lambda_0$, and want to end up in the ground state corresponding to Λ_F: a) at $t = 0$ set $\Lambda = \Lambda_{\text{B}} = \omega_0\omega_F/4 - 1$ (where $\omega_0 = \sqrt{1+\Lambda_0}$ and $\omega_F = \sqrt{1+\Lambda_F}$), b) let the system evolve at fixed Λ_{B} until the time $t = t_{\text{B}} = t_{\text{Rabi}}/(2\sqrt{\omega_0\omega_F})$, and c) change Λ_{B} to Λ_F. Note that the final time is not arbitrary, as before, but is fixed by the initial and final values of Λ. This makes it in practice more unstable with respect to small errors [24]. Under ideal conditions (perfect timing) the method is seen to work well producing fairly low values of ξ_S^2 at times of the order of $t_{\text{Rabi}}/10$ for $N = 100$ atoms with a fidelity of ~ 1 at the final time, see Fig. 4.

In this letter we have presented protocols for fast generation of very coherent-spin-squeezed states in bosonic Josephson junctions. The attained squeezing is the one corresponding to the ordinary adiabatic evolution in the case of repulsive atom-atom interactions, but requires much shorter preparation times. Ordinary adiabatic squeezing is known to improve as N is increased as $\xi_S^2 \propto N^{-1}$ [25], faster than the single-axis-twisting [9] $\xi_S^2 \propto N^{-2/3}$ or the recently proposed production of squeezing in the vicinity of unstable classical points $\xi_S^2 \propto N^{-\sim 0.6}$ [17]. Thus, practical methods of fast-adiabatic driving have important advantages for any future implementation of BJJs where they are used as resources of entanglement/spin squeezing. The present procedures requires a good control of the time variation of the atom-atom scattering length during the desired period, a possibility now at hand in current experimental setups for internal BJJ's. The methods have been obtained by extending recently developed protocols for fast-adiabatic evolution in the case of a single particle evolving in a time dependent Harmonic oscillator potential, to the Bose-Hubbard Hamiltonian. If confirmed experimentally, the present proposal represents a useful step forward in the fast-adiabatic preparation of many-body entangled quantum resources.

The authors thank M. Oberthaler and T. Zibold for useful comments and discussions. This work has been supported by FIS2008-01661, 2009-SGR1289, IT472-10, FIS2009-12773-C02-01, and the UPV/EHU under program UFI 11/55. B. J.-D. is supported by the Ramón y Cajal program. E. T. acknowledges financial support from the Basque Government (Grants No. BFI08.151).

---

[1] M. Albiez, R. Gati, J. Fölling, S. Hunsmann, M. Cristiani, and M. K. Oberthaler, Phys. Rev. Lett. **95**, 010402 (2005).

[2] J. Esteve, C. Gross, A. Weller, S. Giovanazzi, and M. K. Oberthaler, Nature **455**, 1216, (2008).

[3] C. Gross, T. Zibold, E. Nicklas, J. Estève, and M. K. Oberthaler, Nature **464**, 1165 (2010).

[4] T. Zibold, E. Nicklas, C. Gross, and M. K. Oberthaler, Phys. Rev. Lett. **105**, 204101 (2010).

[5] R. Gati and M. K. Oberthaler, J. Phys. B.: At. Mol. Opt. Phys. **40**, R61-R89 (2007).

[6] G. J. Milburn, J. Corney, E. M. Wright, and D. F. Walls, Phys. Rev. A **55**, 4318 (1997).

[7] A. J. Leggett, Rev. Mod. Phys. **73**, 307 (2001).

[8] H.J. Lipkin, N. Meshkov, and A.J. Glick, Nucl. Phys. **62**, 188 (1965).

[9] M. Kitagawa, and M. Ueda, Phys. Rev. A **47**, 5138 (1993).

[10] A. Sorensen, L.-M. Duan, I. Cirac, and P. Zoller, Nature 409, 603 (2001).

[11] J. K. Korbicz, J. I. Cirac, and M. Lewenstein, Phys. Rev. Lett. **95**, 120502 (2005).

[12] X. Chen, A. Ruschhaupt, S. Schmidt, A. del Campo, D. Guéry-Odelin, and J. G. Muga, Phys. Rev. Lett. **104**, 063002 (2010).

[13] X. Chen, and J. G. Muga, Phys. Rev. A **82**, 053403 (2010); X. Chen, E. Torrontegui, and J. G. Muga, Phys. Rev. A **83**, 062116 (2011); E. Torrontegui, S. Ibañez, X. Chen, A. Ruschhaupt, D. Guéry-Odelin and J. G. Muga, Phys. Rev. A. **83**, 013415 (2011).

[14] X. Chen, E. Torrontegui, D. Stefanatos, Jr-Shin Li, and J. G. Muga, Phys. Rev. A **84**, 043415 (2011); E. Torrontegui, X. Chen, M. Modugno, S. Schmidt, A.

Ruschhaupt and J. G. Muga, New J. Phys. **14**, 013031 (2012); A. Ruschhaupt, X. Chen, D. Alonso, and J. G. Muga, arXiv 1206.1691.

[15] V. S. Shchesnovich, and M. Trippenbach, Phys. Rev. A **78**, 023611, (2008).

[16] B. Juliá-Díaz, J. Martorell, and A. Polls, Phys. Rev. A **81**, 063625 (2010).

[17] B. Juliá-Díaz, T. Zibold, M. K. Oberthaler, M. Melé-Messeguer, J. Martorell, and A. Polls, arXiv:1205.6756.

[18] A. V. Turbiner, Commun. Math. Phys. **118**, 467 (1988).

[19] V. V. Ulyanov and O. B. Zaslavskii, Phys. Rep. **216**, 179 (1992).

[20] T. Zibold, private communication.

[21] D. J. Wineland, J. J. Bollinger, W. M. Itano, and F. L. Moore, and D. J. Heinzen, Phys. Rev. A **46**, R6797 (1992).

[22] D. J. Wineland, J. J. Bollinger, W. M. Itano, and D. J. Heinzen, Phys. Rev. A **50**, 67 (1994).

[23] J. Javanainen, and M. Yu. Ivanov, Phys. Rev. A **60**, 2351 (1999).

[24] E. Torrontegui, X. Chen, M. Modugno, A. Ruschhaupt, D. Guéry-Odelin, and J. G. Muga, Phys. Rev. A **85**, 033605 (2012).

[25] C. Gross, arXiv:1203.5359.